

Diapycnal Mixing in a Coastal Regime – AESOP

Michael C. Gregg
Applied Physics Laboratory, University of Washington
1013 NE 40th St.
Seattle, WA 98105-6698
Phone: 206-543-1353 (GreggFax: 206-543-6785
Email: gregg@apl.washington.edu

Grant # N00014-15-1-0362

LONG-TERM GOALS

To identify the major processes producing mixing in the upper ocean and to understand their dynamics sufficiently well to permit accurate parameterization of mixing for use in numerical models.

OBJECTIVES

These measurements during August 2006 were the first attempt we know of to survey a coastal domain with sufficient coverage to assess how mixing levels vary across the domain. Previous measurements have been concentrated in sub-regions, often revealing particular mixing processes, but insufficient to represent mixing throughout a regional model.

APPROACH

We ran lines of microstructure profiles 5-10 km long (Fig. 1), balancing needs for rapid temporal sampling against spatial windows containing at least some structure. Staying with each line for 12.5 hours resolved changes produced by the M2 twice-daily tide, and some lines were rerun at different phases of the monthly tide. As we began to understand patterns of tidal currents and mixing, the original set of lines was modified to reveal pulsing of water in and out of the large canyon splitting the bay down its middle. Powerful Doppler sonars installed on R/V Revelle by Rob Pinkel at Scripps, provided excellent velocity records, supplemented by a 300 kHz ADCP we installed on the bottom the bay's southern half.

WORK COMPLETED

Gregg and Horne (2009) examine mixing in aggregations of fish, probably anchovies, found in the bay. Turbulence was intense, with most dissipation rates at 10^{-6} to 10^{-5} W kg⁻¹, but mixing efficiencies were very low, typically 0.002 compared with 0.2 in turbulent patches produced by shear instabilities. As a result, even though fish schools contributed half of the average dissipation, they made no significant difference in average diapycnal diffusivity. This, however, is a consequence of background diffusivities averaging ten times those found earlier on the New England continental shelf, also during late summer.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2009		2. REPORT TYPE		3. DATES COVERED 00-00-2009 to 00-00-2009	
4. TITLE AND SUBTITLE Diapycnal Mixing in a Coastal Regime - AESOP				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington, Applied Physics Laboratory, 1013 NE 40th St, Seattle, WA, 98105				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			




Fig. 1 MMP (m) and AMP (a) sampling groups in Monterey Bay, overlaid with tidal ellipses. A 300 kHz ADCP was mounted on the bottom at the red triangle. Thin shallow bathymetry contours are at 10 m intervals, and dark contours mark 100 m intervals.

RESULTS

Presently, we are trying to identify other mixing processes in the bay and use that to understand the levels and patterns, then comparing these results with those from other continental shelves. As an example of the patterns, not yet understood, Figure 2 compares average diapycnal diffusivities for MMP groups 10 and 11. Group 10 increases only slightly with depth, whereas group 11 increase much more strongly, reaching $10^{-2} \text{ m}^2 \text{ s}^{-1}$ at the bottom, compared to only one tenth that at the bottom of group 10. These two groups span most of the range of MMP averages.

The major puzzle is why levels are so much higher than on the New England shelf, as overall internal waves are consistent with Levine's (2002) adaptation of Garrett and Munk's (1975) internal waves (Fig. 3). This modification provides a natural reference for internal wave fields in shallow water.




Fig. 2 Average diapycnal diffusivities for mmp groups 10 (left) and 11 (right)




Fig. 3 Clockwise (black) and counter-clockwise frequency spectra at 38 m from the 300 kHz ADCP mounted on the bottom. Levine's spectrum is shown by the green line, and blue lines mark inertial and tidal frequencies.

IMPACT/APPLICATIONS

Gregg and Horne (2009) provide the first observations of turbulence within aggregations of fish, demonstrating that in at least some cases mixing efficiency is much lower than assumed by estimates that biomixing may be a major mixing process.

RELATED PROJECTS

These data, being analyzed in parallel with observations from the Black Sea, Aegean, and Philippines, should bound levels of mixing and processes in coastal regimes, information needed to convince modelers that mixing varies significantly enough to warrant allowing for this in numerical simulations.

MOST SIGNIFICANT ACCOMPLISHMENT

Quantifying mixing intensities and efficiencies with a fish aggregation.

PUBLICATIONS

Gregg, M.C. and J.K. Horne, Turbulence, acoustic backscatter and pelagic nekton in Monterey Bay, *J. Phys. Oceanogr*, 39, 1097-1114, 2009.